

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
11 October 2001 (11.10.2001)

PCT

(10) International Publication Number
WO 01/75161 A2

- (51) International Patent Classification⁷: **C12Q 1/68** (74) Agent: **NEBEL, Heidi, S.**; Zarley, McKee, Thomte, Voorhees & Sease, Suite 3200, 801 Grand Avenue, Des Moines, IA 50309-2721 (US).
- (21) International Application Number: **PCT/US01/10076**
- (22) International Filing Date: **29 March 2001 (29.03.2001)** (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data: **09/538,165** **30 March 2000 (30.03.2000)** **US**
- (71) Applicant (*for all designated States except US*): **IOWA STATE UNIVERSITY RESEARCH FOUNDATION** [US/US]; 310 Lab of Mechanics, Ames, IA 50011 (US).
- (72) Inventors; and
- (73) Inventors/Applicants (*for US only*): **ROTHSCHILD, Max, F.** [US/US]; 310 Lab of Mechanics, Ames, IA 50011-2131 (US). **EMNETT, Rebecca** [US/US]; 310 Lab of Mechanics, Ames, IA 50011 (US). **KIM, Kwan, Suk** [US/US]; 310 Lab of Mechanics, Ames, IA 50011 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— *without international search report and to be republished upon receipt of that report*

[Continued on next page]

(54) Title: **GENETIC MARKERS FOR IMPROVED MEAT CHARACTERISTICS IN ANIMALS**

1 ACAAGAATCT GCATTACCCC ATGTACTTTT TCATCTGTAG CCTGGCTGTG

51 GCTGATATGC TGGTGAGCGT TTCCAATGGG TCAGAAACCA TTGTCATCAC

101 CCTATTAAAC AGCACGGACA CGGACGCACA GAGTTTCACA GTGAATATTG

151 ATAATGTCAT TGA CTCAGTG ATCTGTAGCT CCTTACTCGC CTCAATTTGC

201 AGCCTGCTTT CGATTGCAGT GGACAGGTAT TTTACTATCT TTTATGCTCT

251 CCAGTACCAT AACATTATGA CAGTTAAGCG GGTTGGAATC ATCATCAGTT

301 GTATCTGGGC AGTCTGCACG GTGTCGGGTG TTTTGTTTCAT CATTACTCA

351 GATAGCAGTG CTGTTATTAT CTGCCTCATA ACCGTGTTCT TCACCATGCT

401 GGCTCTCATG GCTTCTCTCT ATGTCCACAT GTTCCTCATG GCCAGACTCC

451 ACATTAAGAG GATCGCCGTC CTCCCAGGCA CTGGCACCAT CCGCCAAGGT

501 GCCAACATGA AGGGGGCAAT TACCCTGACC ATCTTGATTG GGGTCTTTGT

551 GGTCTGCTGG GCCCCCTTCT TCCTCCACTT AATATTCTAT ATCTCCTGCC

(57) Abstract: Genetic markers in the porcine melanocortin-4 receptor (MC4R) gene are disclosed which are associated with favorable meat quality traits including, drip loss, marbling, pH and color. Further, novel sequence data from regions of the gene disclosed which may be used in a PCR test screen for the presence of the marker. The genetic marker may be used to screen animals for breeding purposes which have the desired traits. Kits which take advantage of the PCR test are also disclosed.

WO 01/75161 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE: GENETIC MARKERS FOR IMPROVED MEAT
CHARACTERISTICS IN ANIMALS

FIELD OF THE INVENTION

5 The present invention relates to a method of genetically evaluating animals by assaying for the presence of at least one genetic marker which is indicative of one or more traits associated with meat quality. In particular, the method analyzes for variation in the melanocortin-4 receptor (MC4R) gene or other variations associated therewith which are indicative of these favorable
10 traits.

BACKGROUND OF THE INVENTION

Genetic differences exist among individual animals as well as among breeds which can be exploited by breeding techniques to achieve animals with
15 desirable characteristics. For example, Chinese breeds are known for reaching puberty at an early age and for their large litter size, while American breeds are known for their greater growth rates and leanness. Often, however, heritability for desired traits is low, and standard breeding methods which select individuals based upon phenotypic variations do not take fully into
20 account genetic variability or complex gene interactions which exist.

Restriction fragment length polymorphism (RFLP) analysis has been used by several groups to study pig DNA. Jung et al., Theor. Appl. Genet., 77:271-274 (1989), incorporated herein by reference, discloses the use of RFLP techniques to show genetic variability between two pig breeds. Polymorphism
25 was demonstrated for swine leukocyte antigen (SLA) Class I genes in these breeds. Hoganson et al., Abstract for Annual Meeting of Midwestern Section of the American Society of Animal Science, March 26-28, 1990, incorporated herein by reference, reports on the polymorphism of swine major histocompatibility complex (MHC) genes for Chinese pigs, also demonstrated
30 by RFLP analysis. Jung et al., Theor. Appl. Genet., 77:271-274 (1989), incorporated herein by reference, reports on RFLP analysis of SLA Class I

genes in certain boars. The authors state that the results suggest that there may be an association between swine SLA/MHC Class I genes and production and performance traits. They further state that the use of SLA Class I restriction fragments, as genetic markers, may have potential in the future for
5 improving pig growth performance.

The ability to follow a specific favorable genetic allele involves a novel and lengthy process of the identification of a DNA molecular marker for a major effect gene. The marker may be linked to a single gene with a major effect or linked to a number of genes with additive effects. DNA markers have
10 several advantages; segregation is easy to measure and is unambiguous, and DNA markers are co-dominant, i.e., heterozygous and homozygous animals can be distinctively identified. Once a marker system is established selection decisions could be made very easily, since DNA markers can be assayed any time after a tissue or blood sample can be collected from the individual infant
15 animal, or even an embryo.

The use of genetic differences in receptor genes has become a valuable marker system for selection. For example, United States Patents 5,550,024 and 5,374,526 issued to Rothschild et al. disclose a polymorphism in the pig estrogen receptor gene which is associated with larger litter size, the
20 disclosure of which is incorporated herein by reference. United States number 5,935,784 discloses polymorphic markers in the pig prolactin receptor gene which are associated with larger litter size and overall reproductive efficiency. Perhaps one of the most important characteristics for any meat producing animal is meat quality. Meat quality is a difficult characteristic to assess, as
25 many different aspects, both objective and subjective, make up the overall trait. The list of factors which determine quality in meat, as with other foods, is rather long (Wood et al., Proceedings of The Nutrition Society (1999) 58:363-70). It includes freedom from microbiological hazards (food safety) and prevention of animal exploitation (animal welfare). It also includes the
30 sensory appeal of meat, i.e. its taste or eating quality, and perceived healthiness, especially in relation to the amount and type of fat.

The quality of raw pig meat is influenced by a large number of genetic and non-genetic factors. The latter include farm, transport, slaughter and processing conditions. Meat scientists have performed a substantial amount of research on these factors, which has led to considerable quality improvement.

5 Part of the research has also been dedicated to the genetic background of the pigs, and several studies have revealed the importance of genetic factors. This has made the industry aware that selective breeding of pigs and the use of gene technology can play an important role in enhancing pork quality.

10 Information at DNA level can help to fix a specific major gene, but it can also assist the selection of quantitative trait for which we already select.

Molecular information in addition to phenotypic data can increase the accuracy of selection and therefore the selection response. The size of the extra response in such a Marker Assisted Selection (MAS) program has been considered by many workers from a theoretical point of view. In general
15 terms, MAS is more beneficial for traits with a low heritability and which are expensive to measure phenotypically. Meat quality in particular qualifies as an excellent opportunity to utilize MAS. For example, Meuwissen, T.H.E. and Goddard, M.E.(1996) "The use of Marker Haplotypes in Animal Breeding Schemes", Genet. Sel. Evol., 28 161-176 considered the impact of Marker
20 Assisted Selection for traits such as reproduction and meat quality that are difficult to progress using traditional methods. their results are extremely encouraging, showing that for traits such as meat quality, where the trait is measured after slaughter, an additional response of up to 64% could be achieved.

25 Indeed, the best approach to genetically improve meat quality is to find relevant DNA-markers directly in the population under selection. Meat quality measurements can be performed continuously on some animals from the nucleus populations of breeding organizations. Since a full assessment of meat quality can only be done after slaughter, the data must be collected on
30 culled animals and cannot be obtained on potential breeding animals.

This phenotypic meat quality data is collected in order to enable the detection of relevant DNA markers, and to validate markers from experimental populations or to test candidate genes. Significant markers or genes can then be included directly in the selection process. An advantage of the molecular information is that we can obtain it already at very young age of the breeding animal, which means that animals can be preselected based on DNA markers before the growing performance test is completed. This is a great advantage for the overall testing and selection system.

It can be seen from the foregoing that a need exists for a method for improving meat quality characteristics in animals by identifying and selecting animals with the improved meat characteristics.

An object of the present invention is to provide a genetic marker based on or within the MC4R gene which is indicative of favorable meat characteristics such as those evidenced by pH, marbling, color and drip loss.

Another object of the invention is to provide an assay for determining the presence of this genetic marker.

A further object of the invention is to provide a method of evaluating animals that increases accuracy of selection and breeding methods for the desired traits.

Yet another object of the invention is to provide a PCR amplification test which will greatly expedite the determination of presence of the marker.

An additional object of the invention is to provide a kit for evaluating a sample of animal DNA for the identified genetic marker.

These and other objects, features, and advantages will become apparent after review of the following description and claims of the invention which follow.

SUMMARY OF THE INVENTION

This invention relates to the discovery of a polymorphism within the melanocortin-4 receptor (MC4R) gene which is associated with meat quality traits in animals. This gene is highly conserved among species and it is

expected that the different alleles disclosed herein will also correlate with variability in this gene in other meat producing animals such as bovine, sheep, chicken; etc. This polymorphic site has been previously described in an earlier patent application PCT/US99/16862, publication number WO 00/06777 the disclosure of which is incorporated herein. In the earlier application this site was found to significantly correlate with weight gain and feed intake, in other words, traits involving growth rate of the pig. Surprisingly, as fast growth is generally considered to be negatively correlated with meat quality, the marker has now been shown to correlate with favorable meat characteristics such as pH level, marbling, color, and drip loss. These multigenic characteristics have been previously difficult to associate with quantitative trait loci and current improvements in meat characteristics have centered around understanding and controlling the numerous factors i.e. on a farm, transport, and/or slaughter plant handling influencing meat quality including the incidence of PSE (pale, soft, exudative), RSE (red, soft, exudative), and DFD (dark, firm, dry) meat. According to the invention, the association of the MC4R polymorphism with the these trait(s) enables genetic markers to be identified for specific breeds or genetic lines to identify animals with favorable meat characteristics early in the animal's life.

The marker genotype consists of a polymorphism within the MC4R gene that results in a guanine to adenine transition and a missense mutation of aspartic acid(D) codon (GAU) into asparagine(N) codon (AAU) at a position corresponding to amino acid position 298 of the human MC4R protein resulting in a *TaqI* restriction site in one allele of the gene. In one embodiment of the invention a *TaqI* restriction pattern which identifies the polymorphism is used to assay for the presence or absence of markers associated with the desirable meat traits. The invention includes assays for detection of the marker, or markers linked thereto as well as the sequence characterization of the polymorphism and includes novel sequences in the MC4R gene which may be used to design amplification primers for such an assay (SEQ ID NO:1). Additionally, the invention includes a method for using

the assay in breeding programs for animal selection and a kit for performing the assay.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figure 1 is the sequence listing for MC4R in pigs (SEQ ID NO:1). "X" represents the site of the polymorphism.

 Figure 2 represents a comparison of the DNA sequence between the human (SEQ ID NO:2) and the porcine (SEQ ID NO:1) MC4R gene.

 Figure 3 represents a comparison of the amino acid sequence between
10 the human (SEQ ID NO:3) and the porcine (SEQ ID NO:4) MC4R gene.

 Figures 4a, 4b, and 4c are linkage reports for MC4R from CRI-MAP.

 Figure 5 depicts partial nucleotide and amino acid sequences of the porcine MC4R gene. The amino acid translation shows an amino acid substitution at codon 298.

15 Figure 6 depicts multiple-alignments of the putative seventh transmembrane domain of porcine MC4R with other MCRs and GPCRs. The "*" represents the predicted sequence positions for porcine MC4R (SEQ ID NO:11). The other amino acid sequences were obtained from the GenBank database (accession numbers P32245 (SEQ ID NO:12), P70596 (SEQ ID
20 NO:13), P41983 (SEQ ID NO:14), P56451 (SEQ ID NO:15), P34974 (SEQ ID NO:16), P41968 (SEQ ID NO:17), P33033 (SEQ ID NO:18), Q01718 (SEQ ID NO:19), Q01726 (SEQ ID NO:20), Q28031 (SEQ ID NO:21), AF011466 (SEQ ID NO:22), P21554 (SEQ ID NO:23), P18089 (SEQ ID NO:24), P30680 (SEQ ID NO:25), P47211 (SEQ ID NO:26)). The missense variant in porcine MC4R
25 substituted amino acid N for D in the position marked with an arrow. The Asp (D) residue is highly conserved among MCRs, and the Asn (N) residue is well conserved in most other GPCRs.

DETAILED DESCRIPTION OF THE INVENTION

30 The melanocortin-4 receptor (MC4R) has been shown to be an important mediator of long term weight homeostasis. MC4R antagonists can increase

food intake and body weight during chronic administration. Skuladottir, G.V., et al., "Long term orexigenic effect of a novel melanocortin 4 receptor selective antagonist", British J. of Pharm., 126(1):27-34 (1999).

Lu et al., Nature (Oct 27, 1994), 371 (6500):799-802 suggested that the
5 melanocortin receptors are involved in controlling food intake and energy
balance through studying its antagonism to the *agouti* obesity syndrome.
Huszar et al., Cell (Jan. 10, 1997) 88(1):131-41 found that inactivation of the
melanocortin-4 receptor gene (MC4R) resulted in a maturity onset obesity
syndrome in mice and demonstrated a major role of MC4R protein in the
10 regulation of energy balance related to the *agouti* obesity syndrome. In
addition, the MC4R protein mediates the effects of leptin, one of the important
signaling molecules in energy homeostasis (Seeley et al. 1997).

According to the present invention, a variant or polymorphism in the
MC4R gene has been located, and this genetic variability is associated with
15 phenotypic differences in the porcine meat quality traits as evidenced by pH,
marbling, color and drip loss.

In one embodiment of the invention, an assay is provided for detection of
presence of a desirable genotype in animals. The assay involves assaying the
genomic DNA purified from blood, tissue, semen, or other convenient source of
20 genetic material by the use of primers and standard techniques, such as the
polymerase chain reaction (PCR), then digesting the DNA with a restriction
enzyme (e.g., *Taq I* or other enzyme which cleaves at the same G → A site) so
as to yield gene fragments of varying lengths, and separating at least some of
the fragments from others (e.g., using electrophoresis).

25 The fragments may also be detected by hybridizing with a nucleotide
probe (e.g., radio-labeled cDNA probes) that contains all or at least a portion of
the MC4R gene cDNA sequence to the separated fragments and comparing the
results of the hybridization with assay results for a gene sequence known to
have the marker or a sequence known to not have the marker. Selection and
30 use of probes for detection of MC4R sequences based on the known and
disclosed MC4R sequences is generally known to those skilled in the art. The

probe may be any sequence which will hybridize to the separated digestion products and allow for detection.

Another embodiment of the invention provides a kit for assaying the presence in a MC4R gene sequence of a genetic marker. The marker being
5 indicative of heritable traits of meat quality characteristics. The kit in a preferred embodiment also includes novel PCR primers comprising 4-30 contiguous bases on either side of the polymorphism to provide an amplification system allowing for detection of the G → A Transition polymorphism by PCR digestion of PCR products. The sequence surrounding
10 the polymorphic site is shown in SEQ ID NO:1, Figure 1. Several primers have also been disclosed including SEQ ID NOS:5 and 6, SEQ ID NOS:9 and 10; and mapping primers 7 and 8. The preferred primers are SEQ ID NO:8 and SEQ ID NO:10.

A further embodiment comprises a breeding method whereby an assay
15 of the above type is conducted on a plurality of DNA samples from different animals or animal embryos to be selected from and based on the results, certain animals are either selected or dropped out of the breeding program.

According to the invention, in a preferred embodiment, the polymorphism in the MC4R gene identifiable by the *Taq I* restriction pattern,
20 is disclosed. As is known in the art, restriction patterns are not exact determinants of the size of fragments and are only approximate. When the primers SEQ ID NOS:6 and 7 are used the polymorphism is identifiable by three bands from a *Taq I* digestion of the PCR product, 466, 225, and 76 base pairs (bp) for one homozygous genotype (allele 1); two bands, 542 and 225 bp
25 for another homozygous genotype (allele 2); and four bands for the heterozygous genotype (542, 466, 225, and 76 bp). When the preferred primers are used, SEQ ID NOS:10 and 11 the bands upon *taq* digestion include 156 and 70 bp for allele 1 and one 226bp fragment for allele 2. Those of skill in the art will appreciate that the design of alternate primers PCR conditions and
30 restriction patterns for identifying the presence of allele 2 using the MC4R sequence data herein or other data for closely linked loci represent nothing

more than routine optimization of parameters and are intended to be within the scope of the invention. The marker for improved meat characteristics as evidenced by all four meat quality measurements observed herein (allele 2). The allele 2 genotype was previously associated with faster growth rate. This is surprising because the current state of the art concluded that there is a negative correlation between growth rate and meat quality.

In addition, the polymorphism associated with the pattern has been identified at the nucleotide level. The polymorphic *Taq I* site was sequenced along with the general surrounding area. See SEQ ID NO: 1. The sequences surrounding the polymorphism have facilitated the development of a PCR test in which a primer of about 4-30 contiguous bases taken from the sequence immediately adjacent to the polymorphism is used in connection with a polymerase chain reaction to greatly amplify the region before treatment with the *Taq I* restriction enzyme. The primers need not be the exact complement; substantially equivalent sequences are acceptable.

From sequence data, it was observed that in allele 2 a guanine is substituted with an adenine at position 678 of the PCR product shown in Figure 1 or at position 298 of the analogous human MC4R amino acid of the MC4R protein changing the aspartic acid codon (GAU) into an asparagine codon (AAU). The PCR test for the polymorphism used a forward primer of 5'-TGG CAA TAG CCA AGA ACA AG-3' (SEQ. ID NO: 5) and a reverse primer of 5'-CAG GGG ATA GCA ACA GAT GA-3' (SEQ. ID NO: 6). Pig specific primers used for physical mapping were a forward primer of 5'-TTA AGT GGA GGA AGA AGG-3' (SEQ. ID NO: 7) and a reverse primer of 5'-CAT TAT GAC AGT TAA GCG G-3' (SEQ ID NO:8). The resulting amplified product of about 750 bp, when digested with *Taq I*, results in allelic fragments of 466, 225, and 76 bp (allele 1) or 542 and 225 bp (allele 2). The most preferred primers resulting in either 2 or 1 fragment after *Taq I* digestion are SEQ ID NOS:10 and 11. Allele 1 generates fragments of 156 and 70 base pairs while allele 2 generates a single 226 bp fragment.

The marker may be identified by any method known to one of ordinary skill in the art which identifies the presence or absence of the particular allele or marker, including for example, single-strand conformation polymorphism analysis (SSCP), RFLP analysis, heteroduplex analysis, denaturing gradient gel electrophoresis, allelic PCR, temperature gradient electrophoresis, ligase chain reaction, direct sequencing, minisequencing, nucleic acid hybridization, and micro-array-type detection of the MC4R gene and examination for the polymorphic site. Yet another technique includes an Invader Assay which includes isothermal amplification that relies on a catalytic release of fluorescence. See Third Wave Technology at www.twt.com all of which are intended to be within the scope of the invention.

One or more additional restriction enzymes and/or probes and/or primers can be used. Additional enzymes, constructed probes, and primers can be determined by routine experimentation by those of ordinary skill in the art and are intended to be within the scope of the invention.

Other possible techniques include non-gel systems such as TaqManTM (Perkin Elmer). In this system, oligonucleotide PCR primers are designed that flank the mutation in question and allow PCR amplification of the region. A third oligonucleotide probe is then designed to hybridize to the region containing the base subject to change between different alleles of the gene. This probe is labeled with fluorescent dyes at both the 5' and 3' ends. These dyes are chosen such that while in this proximity to each other the fluorescence of one of them is quenched by the other and cannot be detected. Extension by *Taq* DNA polymerase from the PCR primer positioned 5' on the template relative to the probe leads to the cleavage of the dye attached to the 5' end of the annealed probe through the 5' nuclease activity of the *Taq* DNA polymerase. This removes the quenching effect allowing detection of the fluorescence from the dye at the 3' end of the probe. The discrimination between different DNA sequences arises through the fact that if the hybridization of the probe to the template molecule is not complete, i.e., there is a mismatch of some form, the cleavage of the dye does not take place. Thus,

only if the nucleotide sequence of the oligonucleotide probe is completely complementary to the template molecule to which it is bound will quenching be removed. A reaction mix can contain two different probe sequences each designed against different alleles that might be present, thus, allowing the
5 detection of both alleles in one reaction.

Though the use of RFLPs is one method of detecting the polymorphism, other methods known to one of ordinary skill in the art may be used. Such methods include ones that analyze the polymorphic gene product and detect polymorphisms by detecting the resulting differences in the gene product.

10 Though the preferred method of separating restriction fragments is gel electrophoresis, other alternative methods known to one skilled in the art may be used to separate and determine the size of the restriction fragments.

It is possible to indirectly select for the polymorphism with alternative DNA markers and these methods are also within the scope of the invention.

15 One can establish a linkage between specific alleles of alternative DNA markers and alleles of DNA markers known to be associated with the MC4R gene which have previously been shown to be associated with a particular trait. Examples of markers on the published PiGMap chromosome map which are linked to the MC4R gene include S0331, BHT0433, and S0313. This is
20 also true for other species as well, for example in human the MC4R gene is located at chromosome 18q21.3-q22.

The reagents suitable for applying the methods of the present invention may be packaged into convenient kits. The kits provide the necessary materials, packaged into suitable containers. At a minimum, the kit contains
25 a reagent that identifies the polymorphism in the MC4R gene that is associated with the trait of meat quality. Preferably, the reagent that identifies the polymorphism is a PCR set (a set of primers, DNA polymerase, and four nucleoside triphosphates) that hybridize with the MC4R gene or a fragment thereof. Preferably, the PCR set and restriction enzyme that cleaves
30 the MC4R sequence in at least one place are included in the kit. Preferably, the kit further comprises additional means, such as buffers or reagents, for

detecting or measuring the detectable entity or providing a control. Other reagents used for hybridization, prehybridization, DNA extraction, visualization, and similar purposes may also be included, if desired.

The genetic markers, methods, and kits of the invention are useful in a breeding program to identify and/or to select for meat characteristics in a breed, line, or population of animals. Continuous selection and breeding of animals that are at least heterozygous and preferably homozygous for the desired polymorphism associated with the particular trait would lead to a breed, line, or population having those desired traits. Thus, the marker is a selection tool.

The following examples are offered to illustrate, but not limit the invention.

15

EXAMPLE 1

Melanocortin 4 Receptor PCR-RFLP Test - *TaqI* polymorphism and Genetic Linkage Mapping of MC4R Gene

Primers:

20

Primers were designed from homologous regions of human and rat MC4R sequences (Genbank Accession No. s77415 and u67863, respectively). These primers were used to amplify a 750-bp sequence of the porcine MC4R gene.

25

MC4R1: 5' TGG CAA TAG CCA AGA ACA AG 3' (SEQ ID NO:5)

MC4R4: 5' CAG GGG ATA GCA ACA GAT GA 3' (SEQ ID NO:6)

PCR Conditions:

Mix 1:	10X Promega Buffer	1.0 μ L
	25 mM MgCl ₂	0.6 μ L

dNTPs mix (2.5mM each)	0.5 μ L
25 pmol/ μ L MC4R1	0.1 μ L
25 pmol/ μ L MC4R4	0.1 μ L
dd sterile H ₂ O	7.5 μ L
<i>Taq</i> Polymerase (5 U/ μ L)	0.07 μ L
Genomic DNA (12.5 ng/ μ L)	1.0 μ L

Ten μ L of Mix 1 and DNA were combined in reaction tube, then overlaid with mineral oil. The following PCR program was run: 94°C for 2 min.; 35 cycles of 94°C for 30 sec.; 58°C 1 min., and 72°C 1 min. 30 sec.; followed by a
5 final extension at 72°C for 15 min.

Five μ L of the PCR reaction product was checked on a standard 1% agarose gel to confirm amplification success and clean negative control. Product size is approximately 750 base pairs. Digestion was performed by the following procedure.

10

<u><i>Taq</i>I Digestion Reaction</u>	<u>10 μL reaction</u>
PCR product	5.0 μ L
10X <i>Taq</i> I NE Buffer	1.0 μ L
BSA (10mg/ml)	0.1 μ L
<i>Taq</i> I enzyme (20 U/ μ L)	0.5 μ L
dd sterile H ₂ O	3.4 μ L

15

A cocktail of the buffer, enzyme, BSA, and water was made. Five μ L was added to each reaction tube containing the DNA. The mixture was then incubated at 65°C for at least 4 hours to overnight. Loading dye was mixed with the digestion reaction and the total volume was loaded on a 3% agarose gel. The major bands for allele 1 are about 466, 225, and 76 bp. The allele 2 genotype bands are 542 and 225 bp. The heterozygote genotype has both allele 1 and allele 2.

Results

The amplified PCR product is about 750 bp. The sequence of the PCR product confirmed that the PCR product is MC4R gene with 97.6%, and 92.2% identities at the amino acid and DNA level, respectively, to corresponding human sequences. (see Figs. 2 and 3).

The *TaqI* digestion of the PCR product produced allelic fragments of 466, 225, and 76 bp (allele 1), or 542 and 225 bp (allele 2). The heterozygote genotype has both types of alleles. Mendelian inheritance was observed in three three-generation international reference families, which were used to map this gene by linkage analysis.

The polymorphism between allele 1 and allele 2 resulting from a G → A transition at position 678 of the PCR product revealed a missense mutation of Aspartic acid codon (GAU) into Asparagine codon (AAU) at position 298 amino acid of MC4R protein. (See Figure 1, SEQ ID NO:1).

Allele frequencies were determined by genotyping of DNA samples from a small number of animals from different breeds (Table 1). Allele 1 was observed with a frequency of 1 in Meishan, but was not observed or observed at very low frequency in Hampshire and Yorkshire. The frequencies of allele 1 in Landrace and Chester White were 0.5, respectively.

Figures 2 and 3 illustrate the differences between the DNA and amino acid sequences of the human and porcine MC4R gene (SEQ ID NOS:2-4).

TABLE 1
The Frequency of Allele 1 in Different Pig Breeds

Breed	# Animals	Freq. Allele 1
Meishan	8	1
Large White	8	0.56
Yorkshire	6	0.08
Hampshire	5	0
Landrace	4	0.5
Chester White	4	0.5
Minzu	2	1
Wild Boar	2	1

5 Linkage Analyses

Two-point and multi-point linkage analyses were performed on the genotypes of international reference families. See Figs. 4a-4c. The data were analyzed by using the CRI-MAP program. MC4R was significantly linked to several markers on porcine chromosome (SSC) 1. The most closely linked
 10 markers (recombination fraction and LOD score in parentheses) are SO331 (0.02, 21.97), BHT0433 (0.02, 21.32), and SO313 (0.00, 17.76) by two-point linkage analysis. A multi-point linkage analysis produced the best map order of markers and MC4R (with distance in Kosambi cM): KGF-5.8-CAPN3-2.5-MEF2A-6.1-MC4R-5.6-SO313.

15 Somatic cell hybrid panel of pig and rodent was used to assign MC4R to a cytogenetic region. PCR products from pig specific primers were amplified in clones 7, 8, 16, 18, and 19. MC4R was localized to SSC1q 22-27.

EXAMPLE 2

Discovery of a Missense Variant of the Porcine Melanocortin-4
Receptor (*MC4R*) Gene

5 To determine if there was an association of this *MC4R* polymorphism with phenotypic variation the mutation was tested in a large number of individual animals from several different pig lines. Analyses of growth and performance test records showed significant associations of *MC4R* genotypes with backfat, growth rate and feed intake in a number of lines. It is probable
10 that the variant amino acid residue of the *MC4R* mutation causes a significant change of the MC4R function. These results support the functional significance of a pig *MC4R* missense mutation and suggest that comparative genomics based on model species may be equally important for application to farm animals as they are for human medicine.

15 Identification of mutations in the *leptin* and the *leptin receptor* has provided some information on genetic components involved in the regulation of energy balance (Zhang et al. 1994; Tartaglia et al. 1995). Genetic studies using animal models have facilitated the identification of major genetic causes of obesity (Andersson 1996; Pomp 1997; Giridharan 1998). Furthermore,
20 several other genes involved in the neural signaling pathway of energy homeostasis have been identified (Flier and Maratos-Flier 1998; Schwartz et al. 1999). Of particular interest among candidate signaling molecules involved in the regulation of energy homeostasis is the melanocortin-4 receptor (MC4R). The MC4R response to leptin signaling is a link between food intake and body
25 weight (Seeley et al. 1997; Marsh et al. 1999). Neuropeptide Y (NPY) signaling in the central nervous system is also mediated by the MC4R protein (Kask et al. 1998). Several mutations in MC4R including frameshift and nonsense mutations are associated with dominantly inherited obesity in humans (Vaisse et al. 1998; Yeo et al. 1998). Some other MC4R missense
30 mutations in humans have also been identified (Gotoda et al. 1997; Hinney et

al. 1999) but the functional significance of these mutations has not been characterized.

Selection based on growth characteristics has been of great importance to the pig industry because of costs associated with feeding and consumer preference for lean meat. Efficient genetic improvement in these quantitative traits may be augmented through the use of marker assisted selection (MAS) using high density genetic maps (Dekkers and van Arendonk 1998; Rothschild and Plastow 1999). An important tool in this process is comparative mapping using the well-developed human and mouse gene maps, which assist in the identification of corresponding genomic regions or major genes controlling growth and performance traits in the pig. Biological understanding of complex traits in human or model species offers an alternative approach to identify genes responsible for the traits of economic interest in livestock. Several quantitative trait loci (QTL) linkage scans using phenotypically divergent breeds and candidate gene analyses have been successfully conducted for fatness and growth traits (Yu et al. 1995; Casas-Carrillo et al. 1997; Knorr et al. 1997; Knott et al. 1998; Rohrer et al. 1998; Wang et al. 1998; Paszek et al. 1999), but no individual genes with major effects on growth and performance traits have yet been established for commercial populations. The role of MC4R in feed intake and obesity suggests it may be an important genetic marker for the growth-related traits in the pig.

Materials and Methods

Animals. Pigs were raised under normal production conditions under the care of PIC employees in nucleus farms in the United States and Europe.

PCR amplification of a pig MC4R gene fragment. Primers were designed from homologous regions of human and rat MC4R sequences (GenBank accession no. s77415 and u67863, respectively). The primers were: forward primer: 5'-TGG CAA TAG CCA AGA ACA AG-3' (SEQ. ID NO:5) and reverse primer: 5'-CAG GGG ATA GCA ACA GAT GA-3' (SEQ. ID NO:6). The PCR reaction was performed using 12.5 ng of porcine genomic DNA, 1x PCR

buffer, 1.5 mM MgCl₂, 0.125 mM dNTPs, 0.3 mM of each primer, and 0.35 U *Taq* DNA polymerase (Promega) in a 10 µL final volume. The conditions for PCR were as follows: 2 min at 94°C; 35 cycles of 30 s at 94°C, 1 min at 56°C, 1 min 30 s at 92°C, and a final 15 min extension at 72°C in a Robocycler
5 (Stratagene, La Jolla, CA).

Sequencing and mutation detection. Sequencing of the PCR products from several individual pigs of different breeds was conducted and the sequences were compared to detect any nucleotide change. Sequencing was
10 performed on an ABI sequencer 377 (Applied Biosystems). The porcine MC4R sequence has been submitted to GenBank, and has accession number AF087937. The sequence analysis revealed one nucleotide substitution situated within a *TaqI* restriction enzyme recognition site (Kim et al. 1999). A set of primers was then designed to generate a smaller MC4R gene fragment,
15 which contained only one informative *TaqI* restriction site to specify the polymorphic site and to facilitate the PCR-RFLP test. These primers were: forward 5'-TAC CCT GAC CAT CTT GAT TG-3' (SEQ. ID NO:9) and reverse: 5'-ATA GCA ACA GAT GAT CTC TTT G-3' (SEQ. ID NO:10).

Results

Identification of a missense mutation in the pig MC4R gene. The MC4R
20 gene consists of approximately 1 kb of coding sequence contained within a single exon. About 750 bp of a pig MC4R gene fragment was produced by PCR (Kim et al. 1999). The sequence of the PCR product confirmed that the PCR product is the MC4R gene with 92.2% and 97.6% identities at nucleotide and
25 the amino acid levels, respectively, to the human MC4R sequence. Multiple alignments of the sequences from individual animals of several breeds identified a single nucleotide substitution (G→A; Fig. 5). The polymorphism revealed a missense mutation that replaces aspartic acid (GAU) with asparagine (AAU) at the position identical to amino acid 298 of human MC4R
30 protein. To confirm this base change, we designed pig-specific primers flanking the polymorphic site and analyzed the polymorphism as a *TaqI* PCR-

RFLP gel. Allele 1 produced 156 and 70 bp fragments and allele 2 produced a 226 bp fragment as the PCR-RFLP. The heterozygote has both allele 1 and 2 fragments. Molecular marker (M) and MC4R genotypes are indicated at the top of each lane.

5

The MC4R missense mutation is within a highly conserved region among melanocortin receptors (MCR). The MCR is a subfamily of G-protein coupled receptors (GPCR) containing certain conserved structural elements common to most other GPCRs, but overall amino acid identities between MCR and other GPCRs are low (Tatro 1996). A multiple-alignment of the predicted amino acid sequences of the pig MC4R with MC4R proteins from other species, other MCR proteins, or representative GPCRs showed that the aspartic acid found at position 298 of the seventh transmembrane domain is very highly conserved in the MCR proteins (Fig. 6). It is interesting to note, however, that this position is occupied by asparagine in most other GPCRs. The MCR proteins show 40-80% amino acid identity with each other (Tatro 1996), but the second intracytoplasmic loop and the seventh transmembrane domain are highly conserved among MCR proteins (Gantz et al. 1993). Some of the relationships between MCR structure and function have been discovered by the studies of natural and experimental mutations in humans and mice (Robbins et al. 1993; Valverde et al. 1995; Frandberg et al. 1998). These studies indicate that some mutations in highly conserved regions cause structural changes and alter the function of the receptor. The Asp298Asn substitution mutation could have an effect on the function of the receptor. However, this will require further testing but it is known that change of the homologous residue in MC1R (Asp294His) is associated with fair skin and red hair in humans (Valverde et al. 1995).

EXAMPLE 3

Quantity and quality are descriptive terms of great importance in the meat industry. As the live animal is converted to meat and the meat moves
5 along the line of distribution, from slaughters and processors to retailers and finally to consumers, the factor of quality becomes increasingly more important. Obviously, economic considerations influence the concerns for quantity and quality.

The condition of pale, soft and exudative (PSE) pork and generally very
10 high variability of pork quality was recognized and documented by 1960, and both quality "defects" have been viewed as having less value for further processing and being inferior for consumers. Although an enormous amount of research has been directed at the problem through a half-century of effort, surveys of incidence showed, in pork produced in the U.S.A., that 18% was of
15 inferior quality (PSE) in 1963 and 16% in 1992. Thus, the existence of gene markers associated with both the ability to change the levels of traits (i.g. meat color, water holding capacity, tenderness or marbling) as well as to reduce variation in meat quality characteristics provides excellent opportunities for a dramatic improvement in meat quality. First, gene
20 markers allow for significant steps to be made in the desired direction of quality traits; (e.g. improving technological yield of processed ham and reducing moisture (purge) losses of fresh ham and loins, by selecting against the RN gene in pigs). Secondly, gene markers will help reduce meat quality variation, since we can fix relevant genes in the breeding populations.

25 Meat quality is typically measured in slaughter plants in terms of the pH of the meat, color (using several different instruments and methods e.g. Minolta, [Min]), marbling and drip loss.

For example, the following descriptions of desirable meat quality characteristics are generally accepted by the industry based on their economic
30 value at different segments of the pork supply chain:

Loin Minolta Lightness (L^*): The range of 43-47 units (from darker to lighter color) is acceptable, but L^* of 43 is better; i.e., has higher economic value, in general in this range**

5 Loin Japanese Color Score (JCS): The range of 2.5 – 5.0 units (from lighter to darker color) is acceptable, but JCS of 3-4 is better

Loin Marbling (level of intramuscular fat): Generally, higher marbling is better as it is associated with improved meat eating quality characteristics

10 Loin pHu: (ultimate meat acidity measured 24 hours post-mortem; this attribute is the single most important trait of pork quality); -- The range of 5.50 - 5.80 is desirable, but 5.80 is better as it positively influences the color and (low) purge of the meat

Ham Minolta lightness (L^*) The range of 43-52 units is acceptable, but lower (43) is better

Ham pHu: higher; i.e., 5.80, is better

15 Drip loss or purge: the range of 1%-3% is acceptable, but lower is better

** this may be dependent upon market, for example in Japan darker pork is preferred. Sosnicki, A.A., E.R. Wilson, E.B. Sheiss, A. deVries, 1998 "Is there a cost effective way to produce high quality pork?", Reciprocal Meat Conference
20 Proceedings, Vol. 51.

Results

Table 1

Least square means for different MC4R genotypic classes based on a sample of 1146 animals from six genetic lines (preferred class in bold)

5

Trait	Genotype			
	11	12	22	p value
Loin pHu	5.70	5.70	5.73	<0.01
Ham pHu	5.69	5.69	5.72	<0.07
Ham Min L	48.44	48.39	47.38	<0.03
Drip	2.29	2.43	2.10	<0.07
Loin Marbling	2.17	2.18	2.25	<0.42
Days to 110g*	169.2	168.5	166.4	<0.0001

* The samples size for Days was 2366

Significant effects of marker genotype are identified for ultimate pH (pHu), color (Min) and drip loss and a desirable trend is observed for marbling.

10 The size of the effects observed between genotypes while small are of commercial significance in terms of differences in meat quality. It can be seen from the results in Table 1 that allele 2 is the preferred allele in this sample for all four meat quality measures. Interestingly, this is the preferred allele for growth as reported in WO 00/06777. This is a particularly important
15 finding, as it is somewhat unexpected. In general, there is a negative correlation between growth rate and meat quality. Indeed, there is a general perception that meat quality has decreased as breeders have selected for increased growth rate.

In some situations we might anticipate that the associations between
20 the marker genotype and the traits may differ in direction. This will be the case where the marker utilized here is linked to the polymorphism or gene that is causing the effect. In this situation, the MC4R marker will still have utility, once the association has been identified by experimentation.

The traits measured here are only some of the measures that may be used for determining meat quality. Many others can be used that are correlated to these measures. Thus it will be expected that similar effects will be observed for such economically important traits as water holding and
5 binding capacity, curing and cooking yields and that these traits and that these will also extend to related measures of eating quality such as tenderness, juiciness, flavor and taste. See Sosnicki, supra.

The present invention concerns the identification of significant associations between the MC4R marker genotype and meat quality. It will be
10 realized by those skilled in the art that other gene markers located in this region of the swine genome (swine chromosome 1) will also be suitable for marker assisted selection of these traits.

EXAMPLE 4

15 A total of 257 animals from a Pietrain-based line of pigs were slaughtered and meat quality characteristics determined at the time of slaughter and during post-slaughter handling/conditioning for meat production. MC4R genotypes were determined using methods disclosed herein. Associations between marker genotype and MQ traits were then
20 calculated. The results are depicted in Table 2.

Significant associations were obtained for ultimate pH and color (Minolta L) of the ham. Hams from animals of genotype 2,2 will be preferred in markets which favor these characteristics.

The association can be used to select the parents of slaughter pigs or to
25 improve breeding stock by within line selection. Alternatively, ham processors may chose to purchase pigs of the preferred genotype in order to improve the overall quality of the product (2,2 carcasses will be a better color and will be expected to provide greater yield than those from 1,1 or 1,2 animals). In addition, by selecting a single genotype they will also reduce the variation in
30 product quality due to the different MC4R genotypes.

Table 2

MC4R genotype and Ham pHu and Min L for a Pietrain based line selected to be free of the Halothane gene

Genotype	n	pHu	Min L
11	119	5.72	49.70
12	101	5.73	50.03
22	37	5.80	47.83
p		<0.04	<0.09

The present invention describes an association between MC4R genotype and meat quality characteristics such as pH, color and marbling in the pig. These traits are in turn associated with visual appearance and processing and eating quality characteristics such as tenderness.

These traits also describe meat quality in other species such as beef and lamb. Because of the relatively close evolutionary link between pigs and other meat species it can be predicted that variation in this gene is also likely to be associated with meat quality (MQ) in these other species. Polymorphisms can be identified in the MC4R gene of these species using the same approach set out here and the resulting SNPs used for association analysis.

Having described the invention with reference to particular compositions, theories of effectiveness, and the like, it will be apparent to those of skill in the art that it is not intended that the invention be limited by such illustrative embodiments or mechanisms, and that modifications can be made without departing from the scope or spirit of the invention, as defined by the appended claims. It is intended that all such obvious modifications and variations be included within the scope of the present invention as defined in the appended claims. The claims are meant to cover the claimed components and steps in any sequence which is effective to meet the objectives there intended, unless the context specifically indicates to the contrary.

The following citations are hereby incorporated in their entirety by reference:

- Andersson LB (1996), Ann Med., 28:5-7.
- Arden KC (1990), Cytogenet. Cell Genet. 53:161-165.
- 5 Barinaga M (1996), Science 271:913.
- Bray GA (1978). Physiol. Rev. 59, 719-809.
- Casas-Carillo E (1997), J Anim Sci 75:2047-2053.
- Chajlani V, Biochem. Biophys. Res. Commun., 195, 866-873.
- Chen H (1995), Cell 84, 491-495.
- 10 Chua SC (1996). Science 271, 994-996.
- Chung WK (1996). Genome Res. 6, 431-438.
- Cioffi JA (1996). Nature Med. 2, 585-589.
- Cybulsky MI (1991), Proc. Nat. Acad. Sci. 88:7859-7863.
- 15 Dekkers JCM (1998), Genet Res 71:257-275.
- Flier JS (1998), Cell 92:437-440.
- Frandberg P (1997), Biochem Biophys Res Commun 236:489-492.
- Frankel WN (1996), Nat Genet 14:371-373.
- Gantz (1993), J. Biol. Chem. 268, 15174-15179.
- 20 Gantz (1994), Biochem. Biophys. Res. Comm., 200, 1214-1220.
- Gantz I (1993a), J. Biol. Chem. 268, 8246-8250.
- Giridharan NV (1998), Indian J Med res 108:225-242.
- Gotoda T (1997), Diabetologia 40:976-979.
- Goureau A (1996). Genomics. 36:252-262.
- 25 Griffon N (1994), Biochem. Biophys. Res. Comm., 200, 1007-1014.
- Helm J (1994), J. Anim. Science 72:2764.
- Hinney A (1999), J Clin Endocrinol Metab 84:1483-1486.
- Huszar (1997), Cell 88:131-141.
- Kask A (1998), Biochem Biophys Res Commun 248:245-249.
- 30 Kim KS (2000), J Anim Sci 78:791-792.
- Knorr C (1997), Anim Genet 28:124-128.

- Knott SA (1998), Genetics 149:1069-1680.
- Labbe O (1994), Biochemistry, 33, 4543-4549.
- Lee GH (1996), Nature 379, 632-635.
- Lu Dongsi (1994), Nature:371:799-802
- 5 Marsh DJ (1999), Nat Genet 21:119-122.
- Meuwissen, T.H.E. and Goddard, M.E.(1996) "The use of Marker Haplotypes
in Animal Breeding Schemes", Genet. Sel. Evol., 28 161-176
- Montjoy KG (1992), Science 257, 1248-1251.
- 10 Montjoy KG (1994), Mol. Endocrinol. 8, 1298-1308.
- Paszek A (1996), Proceed. Midwest. ASAS Meeting, p. 24.
- Paszek AA (1999), Mamm Genome 10:117-122.
- Phillips M (1996), Nature Genetics 13:18-19.
- Pomp D (1997), Behav Genet 27:285-306.
- 15 Robbins LS (1993), Cell 72:827-834.
- Robic A (1996), Mamm. Genome 7, 438-445.
- Rohrer GA (1998), J Anim Sci 76:2247-2254.
- Rothschild MF (1996), PNAS 93: 201-205.
- Rothschild MF(1999), AgBiotechNet 1:1-8.
- 20 Schwartz MW (1999), Am J Clin Nutr 69:584-96.
- Seeley RJ (1997), Nature 390:349.
- Tartaglia LA (1995), Cell 83:1263-1271.
- Tatro JB (1996), Neuroimmunomodulation 3:259-284.
- Truett GE (1995), Mamm. Genome 6, 25-30.
- 25 Vaisse C (1998), Nat Genet 20:113-114.
- Valverde P (1995), Nat Genet 11:328-330.
- Wang L (1998), J Anim Sci 76:2560-2567.
- Warner CM (1991), Immunogenetics of the MHC, VCH Publishers, NY, NY,
pp. 368-397.
- 30 Winick JD (1996), Genomics 36, 221-222.
- Xy Y (1994), Biol. Reprod. 51:695-699.

- Yeo GS (1998), Nat Genet 20:111-112.
- Yerle M (1996), Cytogenet. Cell Genet. 73, 194-202.
- Youngs CR (1993), J. Anim. Sci., 1561-1565.
- Yu TP (1995), J Anim Sci 73:1282-1288.
- 5 Zhang Y (1994), Nature 372:425-432.
- Ziang ZH (1999), Mamm Genome 12:191-193.

What is claimed is:

1. A method of identifying an animal which possesses a genotype indicative of favorable meat quality traits, said method comprising:
 - a) obtaining a nucleic acid sample from said animal, and
 - b) assaying for the presence of a polymorphism in the MC4R gene of the sample, or a polymorphism linked thereto, said polymorphism being one which is associated with favorable meat quality characteristics such pH, marbling, color and drip loss.
2. The method of claim 1 wherein said polymorphism is characterized by an aspartic acid codon (GAU) which is changed to an asparagine codon (AAU) at amino acid position analogous to amino acid 298 of a human MC4R gene product.
3. The method of claim 1 wherein said aspartic acid codon at position 298 of the MC4R gene product is associated with improved meat characteristics.
4. The method of claim 1 wherein the animal is a pig.
5. The method of claim 1 wherein the step of identifying the polymorphism is a method employing allele specific oligonucleotides.
6. The method of claim 1 wherein said polymorphism is identified by PCR amplification and restriction.
7. The method of claim 1 wherein the step of identifying the polymorphism is selected from the group consisting of restriction fragment length polymorphism (RFLP) analysis, heteroduplex analysis, single strand conformational polymorphism (SSCP) analysis, denaturing gradient gel

electrophoresis (DGGE), temperature gradient gel electrophoresis (TGGE), and use of linked genetic markers.

8. The method of claim 7 wherein the step of identifying the polymorphism
5 comprises RFLP analysis.

9. The method of claim 1 further comprising the step of:
amplifying the MC4R gene sequence.

10 10. The method of claim 9 further comprising the step of
digesting the amplified region with the restriction endonuclease *Taq I*.

11. The amplified gene sequence of claim 10 wherein primers used in the
amplification are selected from the group consisting of SEQ. ID NO:6, SEQ. ID
15 NO:7, SEQ. ID NO:8, SEQ. ID NO:9, SEQ. ID NO:10, and SEQ. ID NO:11.

12. A single strand of an oligonucleotide primer useful for detecting
nucleotide 678 of SEQ ID NO:1, the primer consisting of a nucleotide sequence
having about 4-30 contiguous bases from SEQ ID NO:1 and flanking position
20 678.

13. The oligonucleotide of claim 12 wherein the oligonucleotide has the
nucleotide sequence represented by SEQ ID NO:6.

25 14. The oligonucleotide of claim 12 wherein the oligonucleotide has the
nucleotide sequence represented by SEQ ID NO:7.

15. The oligonucleotide of claim 12 wherein the oligonucleotide has the
nucleotide sequence represented by SEQ ID NO:8.

30

16. The oligonucleotide of claim 12 wherein the oligonucleotide has the nucleotide sequence represented by SEQ ID NO:9.
17. The oligonucleotide of claim 12 wherein the oligonucleotide has the nucleotide sequence represented by SEQ ID NO:10.
18. The oligonucleotide of claim 12 wherein the oligonucleotide has the nucleotide sequence represented by SEQ ID NO:11.
19. The method of claim 1 wherein said primer is SEQ ID NOS: 6 and 7 and wherein said polymorphism is position 678 of said amplified sequence.
20. A method of identifying an animal which possess a desired genotype indicative of favorable meat quality characteristics, the method comprising:
- a) obtaining a nucleic acid sample,
 - b) digesting the sample with a restriction enzyme that recognizes the same site as *Taq I* to obtain fragments,
 - c) separating the fragments obtained from the digestion, and
 - d) identifying the presence or absence of a *Taq I* site at the codon for position for the amino acid codon at position 298 of the MC4R gene product.
21. The method of claim 20 further comprising the step of selecting animals with the desired genotype for breeding.
22. The method of claim 20 wherein the site is identifiable by fragments of 466, 225, and 76 bp when a guanine is present at base 678 and fragments of 542 and 225 bp when an adenine is present when a restriction enzyme which cuts at the same recognition site as *Taq I* is used.

23. The method of claim 20 wherein the site is identifiable by fragments of 156 and 70 bp when allele one is present and 226 bp when allele 2 is present, when a restriction enzyme that acts at the same recognition site as *TaqI* is used.

5

24. The method of claim 20 wherein the step of identifying comprises: detecting the *Taq I* site by amplification.

25. A kit for evaluating a nucleic acid sample from an animal comprising:
10 a reagent in a container that identifies a polymorphism in a MC4R gene.

26. The kit of claim 25 wherein the reagent is a primer that amplifies the MC4R gene or a fragment thereof.

15 27. The kit of claim 25 further comprising: a DNA polymerase which cleaves the MC4R gene, a forward primer, and a reverse primer, wherein the primers are capable of amplifying a region of the MC4R gene which contains a polymorphic site.

20 28. A primer for assaying the presence of a polymorphic *TaqI* site in the MC4R gene wherein the primer comprises a sequence selected from the group consisting of SEQ. ID NO:6, SEQ. ID NO:7, SEQ. ID NO:8, SEQ. ID NO:9, SEQ. ID NO:10, and SEQ. ID NO:11.

25 29. A method for selecting animals for the desired traits of improved meat quality comprising the steps of:

- a) obtaining a nucleic acid sample from an animal,
- b) identifying a polymorphism characterized by a nucleotide position 678 of SEQ ID NO:1 from the MC4R gene, and
- 30 c) selecting the animals which have the nucleotide associated with the desired traits in position 678.

30. A method for an indirect selection for a polymorphism in MC4R wherein specific alleles of an alternative DNA marker are used to make the indirect selection wherein the alternative DNA marker is a linked marker near MC4R.

5 31. The method of claim 30 wherein the linked marker is selected from the group consisting of S0331, BHT0433, and S0313.

32. A method of identifying animals which possess a desired genotype indicative of favorable meat quality traits, the method comprising:
10 determining an association between a MC4R genotype and a trait of interest by obtaining a sample of animals from a line or breed of interest, preparing a nucleic acid sample from each animal in the sample, determining the genotype of the MC4R gene, and calculating the association between the MC4R
genotype and the trait.

15 33. A method of selecting animals which possess a desired MC4R genotype indicative of favorable meat quality traits, the method comprising: obtaining a nucleic acid sample from an animal, identifying the genotype of the MC4R gene of the animal, and selecting those animals which have the genotype
20 associated with the desired traits.

34. A method of determining the potential meat quality (value) of an animal, said method comprising: obtaining a nucleic acid sample from said animal and the assaying for the presence of a polymorphism in the MC4R gene
25 of the sample, or a polymorphism linked thereto, said polymorphism being one which is associated with favorable meat quality characteristics such as pH, marbling, color and drip loss.

35. A method of selecting animals for breeding, said method comprising:
30 a) obtaining a nucleic acid sample from said animal;

- b) assaying for the presence of a polymorphism in the MC4R gene of said sample or a polymorphism linked thereto, said polymorphism being one which is associated with favorable meat quality characteristics such as pH, marbling, color and drop loss; and
- 5 c) using the MC4R genotype as part of a selection index based on the estimated value of the effect.

36. A method of segregating animals in order to prove uniformity at slaughter comprising:

- 10 a) obtaining a nucleic acid sample from said animal; and
- b) assaying for the presence of a polymorphism in the MC4R gene of said sample or a polymorphism linked thereto, said polymorphism being one which is associated with favorable meat quality characteristics such as pH, marbling, color and drop loss.

15

37. A method for selecting animals for breeding, said method comprising:

- a) obtaining a nucleic acid sample from said animal;
- b) assaying for the presence of a polymorphism in the MC4R gene of said sample or a polymorphism linked thereto, said polymorphism
20 being one which is associated with favorable meat quality characteristics such as pH, marbling, color and drop loss; and
- c) selecting animals with a favorable allele for inclusion in breeding stock.

25 38. The method of claim 1 wherein said polymorphism is a single nucleotide polymorphism within the MC4R gene or linked to the MC4R gene such that it is associated with variation in meat quality.

39. The method claim 1 wherein said polymorphism is a complex
30 polymorphism including repeat link variance (e.g., micro-satellites), insertions or deletions.

1 ACAAGAATCT GCATTCAACC ATGTACTTTT TCATCTGTAG CCTGGCTGTG
51 GCTGATATGC TGGTGAGCGT TTCCAATGGG TCAGAAAACCA TTGTCAATCAC
101 CCTATTAAAC AGCACGGACA CGGACGCACA GAGTTTCACA GTGAATATTG
151 ATAAATGTCAT TGAATCAGTG ATCTGTAGCT CCTTACTCGC CTCGAATTTGC
201 AGCCTGCTTT CGATTGCAGT GGACAGGTAT TTTACTATCT TTTATGCTCT
251 CCAGTACCAT AACATTATGA CAGTTAAGCG GGTGGAATC ATCATCAGTT
301 GTATCTGGGC AGTCTGCACG GTGTCGGGTG TTTTGTTCAT CATTACTCA
351 GATAGCAGTG CTGTTATTAT CTGCCCTCATA ACCGTGTTCT TCACCATGCT
401 GGCTCTCATG GCTTCTCTCT ATGTCCACAT GTTCCTCATG GCCAGACTCC
451 ACATTAAGAG GATCGCCGTC CTCCCAGGCA CTGGCACCAT CCGCCAAGGT
501 GCCAACATGA AGGGGGCAAT TACCCTGACC ATCTTGATTG GGTCTTTGT
551 GGCTCTGCTGG GCCCCCTTCT TCCTCCACTT AATATTCTAT ATCTCCTGCC

Fig. 1

601 CCCAGAAATCC ATACTGTGTG TGCTTCAATG CTCACATTAA TTTGTATCTC
651 ATCCTGATCA TGTGTAATTC CATCATCXAT CCCCTGATTT ATGCACTCCG
701 GAGCCAAGAA CTGAGGAAAA CCTTCAAAGA GATCATCTGT TGCTAT

Fig. 1A

```
con-mc4r.seq          10      20      30
                      ACAAAGAATCTGCATTCAACCCATGTACTTTT
                      |||||
s77415                ATATCTTAGTGATTGTGGCAATAGCCAAGAAATCTGCATTCAACCCATGTACTTTT
580 590 600 610 620 630

con-mc4r.seq          40      50      60      70      80      90
                      TCATCTGTAGCCCTGGCTGTGGCTGATATGCTGGTGAGCGTTTCCAATGGGTCAGAAACCA
                      |||||
s77415                TCATCTGCAGCTTGGCTGTGGCTGATATGCTGGTGAGCGTTTCAAATGGATCAGAAACCA
640 650 660 670 680 690

con-mc4r.seq          100     110     120     130     140     150
                      TTGTCAATCACCCCTATTAAACAGCACGGACACGGACGACAGAGTTTCAAGTGAATATTG
                      |||||
s77415                TTATCAATCACCCCTATTAAACAGTACAGATACGGATGCACAGAGTTTCAAGTGAATATTG
700 710 720 730 740 750

con-mc4r.seq          160     170     180     190     200     210
                      ATAATGTCATTGACTCAGTGATCTGTAGCTCCTTACTCGCCTCAATTTCAGCCCTGCTTTT
                      |||||
s77415                ATAATGTCATTGACTCGGTGATCTGTAGCTCCTTGTCTTGATCCATTTCAGCCCTGCTTT
760 770 780 790 800 810
```

Fig. 2A

```
con-mc4r.seq      220      230      240      250      260      270
CGATTGCAGTGGACAGGTATTTTACTATCTTTTATGCTCTCCAGTACCATAACATTATGA
|||||
s77415            CAATTGCAGTGGACAGGTACTTTACTATCTTCTATGCTCTCCAGTACCATAACATTATGA
820      830      840      850      860      870

con-mc4r.seq      280      290      300      310      320      330
CAGTTAAGCGGGTTGGAATCATCATCAGTTGTATCTGGGCAGTCTGCACGGTGTGCGGTG
|||||
s77415            CAGTTAAGCGGGTTGGGATCAGCATAAAGTTGTATCTGGGCAGCTTGACGGTTTCAGGCA
880      890      900      910      920      930

con-mc4r.seq      340      350      360      370      380      390
TTTTGTTCATCATTTTACTCAGATAGCAGTGTGTTATTAATCTGCCATCATAACCGTGTCT
|||||
s77415            TTTTGTTCATCATTTTACTCAGATAGTAGTGTGTCATCATCTGCCCTCATCACCATGTTCT
940      950      960      970      980      990

con-mc4r.seq      400      410      420      430      440      450
TCACCATGTGGCTCTCATGGCTTCTCTATGTCCACATGTTCCCTCATGGCCAGACTCC
|||||
s77415            TCACCATGTGGCTCTCATGGCTTCTCTATGTCCACATGTTCCCTGATGGCCAGGCTTC
1000     1010     1020     1030     1040     1050

con-mc4r.seq      460      470      480      490      500      510
ACATTAAAGAGGATCGCCGTCCTCCAGGCACCTGGCACCATCCGCCAAGGTGCCAACATGA
|||||
s77415            ACATTAAAGAGGATTGCTGTCTCTCCCGGCACCTGGTGCCATCCGCCAAGGTGCCAATATGA
1060     1070     1080     1090     1100     1110
```

Fig. 2B

	520	530	540	550	560	570
con-mc4r.seq	AGGGGGCAATTACCCCTGACCATCTTGATTGGGGTCTTTGGTGGTCTGCTGGGCCCCCTTCT					
	AGGGAGCGGATTACCTTGACCATCCTGATTGGCGTCTTTGTTGTCTGCTGGGCCCCATTCT					
s77415	1120 1130 1140 1150 1160 1170					
	580	590	600	610	620	630
con-mc4r.seq	TCCTCCACTTAATATCTATATCTCTGCCCCAGAAATCCATACTGTGTGCTTCATGT					
	TCCTCCACTTAATATCTACATCTCTTGCTCCAGAAATCCATAATTTGTGTGCTTCATGT					
s77415	1180 1190 1200 1210 1220 1230					
	640	650	660	670	680	690
con-mc4r.seq	CTCAGTTTAATTGTATCTCATCTGATCATGTGTAAATCCATCATCAATCCCCTGATTT					
	CTCAGTTTAACCTTGTATCTCATACTGATCATGTGTAAATCAATCATCGATCCTCTGATTT					
s77415	1240 1250 1260 1270 1280 1290					
	700	710	720	730	740	
con-mc4r.seq	ATGCACTCCGGAGCCAGAACTGAGGAAACCTTCAAAGAGATCATCTGTTGCTAT					
	ATGCACTCCGGAGTCAAGAACTGAGGAAACCTTCAAAGAGATCATCTGTTGCTATCCCC					
s77415	1300 1310 1320 1330 1340 1350					
	750	760	770	780	790	800
s77415	TGGGAGGCCCTTTGTGACTTGTCTAGAGATATTAATGGGGACAGACGCAATATAGG					
	1360 1370 1380 1390 1400 1410					

Fig. 2C

S0082	MC4R	rec. fracs.=	0.05,	lods =	14.74
CGA	MC4R	rec. fracs.=	0.14,	lods =	6.88
S0020	MC4R	rec. fracs.=	0.18,	lods =	5.32
S0079	MC4R	rec. fracs.=	0.12,	lods =	10.35
S0155	MC4R	rec. fracs.=	0.14,	lods =	7.68
S0122	MC4R	rec. fracs.=	0.18,	lods =	5.17
S0313	MC4R	rec. fracs.=	0.00,	lods =	17.76
S0312	MC4R	rec. fracs.=	0.20,	lods =	5.60
S0311	MC4R	rec. fracs.=	0.17,	lods =	7.18
S0416	MC4R	rec. fracs.=	0.20,	lods =	3.21
S0331	MC4R	rec. fracs.=	0.02,	lods =	21.91
S0396	MC4R	rec. fracs.=	0.16,	lods =	7.85
BHT0433	MC4R	rec. fracs.=	0.02,	lods =	21.32
S0536	MC4R	rec. fracs.=	0.03,	lods =	15.61
CAPN3	MC4R	rec. fracs.=	0.12,	lods =	6.65

Fig. 4A

KGF	MC4R	rec.	fracs.=	0.09,	lods =	6.46
MEF2A	MC4R	rec.	fracs.=	0.05,	lods =	14.36
MC4R	MC4R	rec.	fracs.=	0.00,	lods =	26.19
S0082	MC4R	rec.	fracs.=	0.00	0.09,	lods = 15.86
CGA	MC4R	rec.	fracs.=	0.07	0.22,	lods = 7.46
S0020	MC4R	rec.	fracs.=	0.00	0.25,	lods = 6.33
S0079	MC4R	rec.	fracs.=	0.00	0.19,	lods = 11.48
S0155	MC4R	rec.	fracs.=	0.00	0.24,	lods = 9.98
S0122	MC4R	rec.	fracs.=	0.00	0.27,	lods = 7.10
S0313	MC4R	rec.	fracs.=	0.00	0.00,	lods = 17.76
S0312	MC4R	rec.	fracs.=	0.04	0.29,	lods = 7.45

Fig. 4B

S0311	MC4R	rec.	fracs.=	0.00	0.28,	lods =	9.02
S0416	MC4R	rec.	fracs.=	0.00	0.31,	lods =	4.17
S0331	MC4R	rec.	fracs.=	0.05	0.00,	lods =	22.14
S0396	MC4R	rec.	fracs.=	0.03	0.24,	lods =	9.33
BHT0385	MC4R	rec.	fracs.=	0.14	0.36,	lods =	3.46
BHT0433	MC4R	rec.	fracs.=	0.05	0.00,	lods =	21.82
S0536	MC4R	rec.	fracs.=	0.00	0.05,	lods =	15.77
CAPN3	MC4R	rec.	fracs.=	0.00	0.18,	lods =	7.35
KGF	MC4R	rec.	fracs.=	0.00	0.17,	lods =	6.74
MEF2A	MC4R	rec.	fracs.=	0.10	0.00,	lods =	14.52
MC4R	MC4R	rec.	fracs.=	0.00	0.00,	lods =	26.19

Fig. 4C

0	ESR			0.0
		0.18	18.4	
1	S0008			18.4
		0.12	11.9	
7	CGA			30.3
		0.03	2.8	
3	S0312			33.1
		0.05	4.9	
4	S0122			38.1
		0.09	9.4	
8	KGF			47.4
		0.06	5.8	
6	CAPN3			53.2
		0.02	2.5	
9	MEF2A			55.7
		0.06	6.1	
5	MC4R			61.8
		0.06	5.6	
10	S0313			67.4
		0.00	0.0	
11	S0082			67.4
		0.03	3.4	
12	S0079			70.8
		0.03	2.5	
14	S0142			73.3
		0.01	1.0	
13	S0020			74.4
		0.04	4.3	
15	S0311			78.7
		0.00	0.0	
16	S0155			78.7
		0.12	12.2	
17	S0113			90.9
		0.20	21.0	
18	S0302			111.9
		0.22	23.4	
19	S0112			135.3

Fig. 4D





Fig. 5



pMC4RMSHFNLYLILIMCNSIIDPLIYAL.....*
hMC4RMSHFNLYLILIMCNSIIDPLIYAL..... 304
rMC4RMSHFNLYLILIMCNAVIDPLIYAL..... 304
sheep MC5RMSHFNMYLILIMCNSVIDPLIYA..... 286
bovine MC5RMSHFNMYLILIMCNSVIDPLIYA..... 286
bovine MC2RMSLFQVNGVLIMCNAIIDPFIYAL..... 268
hMC3RAHFNTYLVLIMCNSVIDPLIYA..... 327
mMC3RAHFNTYLVLIMCNSVIDPLIYA..... 290
hMC2RMSHFNMYLILIMCNSVMDPLIYA..... 268
hMC1RSYFNLFLLILICNSVVDPLIYA..... 299
bEDG-2RLAYEKFFLLLAEFNSAMNPIIYSYR.... 314
hEDG-4RFLLLAEANSLVNAAVYSCR.... 298
human cannabVFAFCSMLCCLLNSTVNPLIYAL..... 399
hH2ABFQFFFWIGYCNSSLNPVIYTI..... 290
rSSR2FDFVVILTYANSCANPILYAFL.... 315
hGAL1-RLAYSNSSVNPIIYAFL.... 306

Fig. 6

SEQUENCE LISTING

<110> Iowa State University Research Foundation

<120> Genetic Markers for Improved Meat Characteristics in Animals

<130> ISURF 2697

<150> 09/538,165

<151> 2000-03-30

<160> 26

<170> PatentIn version 3.0

<210> 1

<211> 746

<212> DNA

<213> Sus scrofa

<400> 1

```

acaagaatct gcattcaccc atgtactttt tcactctgtag cctggctgtg gctgatatgc      60
tggtgagcgt ttccaatggg tcagaaacca ttgtcatcac cctattaaac agcacggaca      120
cggacgcaca gagtttcaca gtgaatattg ataatgtcat tgactcagtg atctgtagct      180
ccttactcgc ctcaatttgc agcctgcttt cgattgcagt ggacaggat tttactatct      240
tttatgctct ccagtaccat aacattatga cagttaagcg ggttgaatc atcatcagtt      300
gtatctgggc agtctgcacg gtgtcgggtg ttttgttcat catttactca gatagcagt      360
ctgttattat ctgcctcata accgtgttct tcaccatgct ggctctcatg gcttctctct      420
atgtccacat gttcctcatg gccagactcc acattaagag gatcgccgtc ctcccaggca      480
ctggcaccat ccgccaaggt gccaacatga agggggcaat taccctgacc atcttgattg      540
gggtctttgt ggtctgctgg gcccccttct tcctccactt aatattctat atctcctgcc      600
cccagaatcc atactgtgtg tgcttcatgt ctcactttaa tttgtatctc atcctgatca      660
tgtgtaattc catcatcrat cccctgattt atgcactccg gagccaagaa ctgaggaaaa      720
ccttcaaaga gatcatctgt tgctat                                           746

```

<210> 2

<211> 840

<212> DNA

<213> Homo sapiens

<400> 2

```

atatcttagt gattgtggca atagccaaga acaagaatct gcattcaccc atgtactttt      60
tcactctcag cttggctgtg gctgatatgc tggtgagcgt ttcaaatgga tcagaaacca      120
ttatcatcac cctattaaac agtacagata cggatgcaca gagtttcaca gtgaatattg      180

```

ataatgtcat tgactcgggtg atctgtagct ccttgcttgc atccatttgc agcctgcttt 240
 caattgcagt ggacaggtac ttactatct tctatgctct ccagtaccat aacattatga 300
 cagttaagcg ggttggggtc agcataagtt gtatctgggc agcttgacg gtttcaggca 360
 ttttgttcat catttactca gatagtagtg ctgtcatcat ctgcctcatc accatgttct 420
 tcaccatgct ggctctcatg gcttctctct atgtccacat gttcctgatg gccaggcttc 480
 acattaagag gattgctgtc ctccccggca ctggtgccat ccgccaaagg gccaatatga 540
 agggagcgat taccttgacc atcctgattg gcgtctttgt tgtctgctgg gcccattct 600
 tctccactt aatattctac atctctgtc ctccagaatcc atattgtgtg tgcttcatgt 660
 ctcaactttaa cttgtatctc atactgatca tgtgtaattc aatcatcgat cctctgattt 720
 atgcactccg gagtcaagaa ctgaggaaaa ccttcaaaga gatcatctgt tgctatcccc 780
 tgggaggcct ttgtgacttg tctagcagat attaaatggg gacagagcac gcaatatagg 840

<210> 3
 <211> 311
 <212> PRT
 <213> Homo sapiens

<220>
 <221> variant
 <222> 298
 <223> "Xaa" can be any amino acid

<400> 3

Gln Leu Phe Val Ser Pro Glu Val Phe Val Thr Leu Gly Val Ile Ser
 1 5 10 15
 Leu Leu Glu Asn Ile Leu Val Ile Val Ala Ile Ala Lys Asn Lys Asn
 20 25 30
 Leu His Ser Pro Met Tyr Phe Phe Ile Cys Ser Leu Ala Val Ala Asp
 35 40 45
 Met Leu Val Ser Val Ser Asn Gly Ser Glu Thr Ile Ile Ile Thr Leu
 50 55 60
 Leu Asn Ser Thr Asp Thr Asp Ala Gln Ser Phe Thr Val Asn Ile Asp
 65 70 75 80
 Asn Val Ile Asp Ser Val Ile Cys Ser Ser Leu Leu Ala Ser Ile Cys
 85 90 95
 Ser Leu Leu Ser Ile Ala Val Asp Arg Tyr Phe Thr Ile Phe Tyr Ala
 100 105 110
 Leu Gln Tyr His Asn Ile Met Thr Val Lys Arg Val Gly Ile Ser Ile

115	120	125
Ser Cys Ile Trp Ala Ala Cys Thr Val Ser Gly Ile Leu Phe Ile Ile		
130	135	140
Tyr Ser Asp Ser Ser Ala Val Ile Ile Cys Leu Ile Thr Met Phe Phe		
145	150	155 160
Thr Met Leu Ala Leu Met Ala Ser Leu Tyr Val His Met Phe Leu Met		
	165	170 175
Ala Arg Leu His Ile Lys Arg Ile Ala Val Leu Pro Gly Thr Gly Ala		
	180	185 190
Ile Arg Gln Gly Ala Asn Met Lys Gly Ala Ile Thr Leu Thr Ile Leu		
	195	200 205
Ile Gly Val Phe Val Val Cys Trp Ala Pro Phe Phe Leu His Leu Ile		
	210	215 220
Phe Tyr Ile Ser Cys Pro Gln Asn Pro Tyr Cys Val Cys Phe Met Ser		
225	230	235 240
His Phe Asn Leu Tyr Leu Ile Leu Ile Met Cys Asn Ser Ile Ile Asp		
	245	250 255
Pro Leu Ile Tyr Ala Leu Arg Ser Gln Glu Leu Arg Lys Thr Phe Lys		
	260	265 270
Glu Ile Ile Cys Cys Tyr Pro Leu Gly Gly Leu Cys Asp Leu Ser Ser		
	275	280 285
Arg Tyr Ala Pro Pro Glu Asn Asp Ile Xaa Val Ile Cys Asn Phe Ile		
	290	295 300
Asp Glu Asn Thr Ile Ala Leu		
305	310	

<210> 4
 <211> 248
 <212> PRT
 <213> Homo sapiens

<400> 4

Lys Asn Leu His Ser Pro Met Tyr Phe Phe Ile Cys Ser Leu Ala Val	
1	5 10 15
Ala Asp Met Leu Val Ser Val Ser Asn Gly Ser Glu Thr Ile Val Ile	
	20 25 30
Thr Leu Leu Asn Ser Thr Asp Thr Asp Ala Gln Ser Phe Thr Val Asn	
	35 40 45
Ile Asp Asn Val Ile Asp Ser Val Ile Cys Ser Ser Leu Leu Ala Ser	
	50 55 60
Ile Cys Ser Leu Leu Ser Ile Ala Val Asp Arg Tyr Phe Thr Ile Phe	
65	70 75 80

Tyr Ala Leu Gln Tyr His Asn Ile Met Thr Val Lys Arg Val Gly Ile
 85 90 95
 Ile Ile Ser Cys Ile Trp Ala Val Cys Thr Val Ser Gly Val Leu Phe
 100 105 110
 Ile Ile Tyr Ser Asp Ser Ser Ala Val Ile Ile Cys Leu Ile Thr Val
 115 120 125
 Phe Phe Thr Met Leu Ala Leu Met Ala Ser Leu Tyr Val His Met Phe
 130 135 140
 Leu Met Ala Arg Leu His Ile Lys Arg Ile Ala Val Leu Pro Gly Thr
 145 150 155 160
 Gly Thr Ile Arg Gln Gly Ala Asn Met Lys Gly Ala Ile Thr Leu Thr
 165 170 175
 Ile Leu Ile Gly Val Phe Val Val Cys Trp Ala Pro Phe Phe Leu His
 180 185 190
 Leu Ile Phe Tyr Ile Ser Cys Pro Gln Asn Pro Tyr Cys Val Cys Phe
 195 200 205
 Met Ser His Phe Asn Leu Tyr Leu Ile Leu Ile Met Cys Asn Ser Ile
 210 215 220
 Ile Asn Pro Leu Ile Tyr Ala Leu Arg Ser Gln Glu Leu Arg Lys Thr
 225 230 235 240
 Phe Lys Glu Ile Ile Cys Cys Tyr
 245

<210> 5
 <211> 20
 <212> DNA
 <213> Sus scrofa

<400> 5
 tggcaatagc caagaacaag

20

<210> 6
 <211> 20
 <212> DNA
 <213> Sus scrofa

<400> 6
 caggggatag caacagatga

20

<210> 7
 <211> 18
 <212> DNA
 <213> Sus scrofa

<400> 7
 ttaagtggag gaagaagg

18

<210> 8
 <211> 19
 <212> DNA
 <213> Sus scrofa

<400> 8
 cattatgaca gttaagcgg 19

<210> 9
 <211> 20
 <212> DNA
 <213> Sus scrofa

<400> 9
 taccctgacc atcttgattg 20

<210> 10
 <211> 22
 <212> DNA
 <213> Sus scrofa

<400> 10
 atagcaacag atgatctctt tg 22

<210> 11
 <211> 24
 <212> PRT
 <213> Sus scrofa

<400> 11

Met Ser His Phe Asn Leu Tyr Leu Ile Leu Ile Met Cys Asn Ser Ile
 1 5 10 15

Ile Asp Pro Leu Ile Tyr Ala Leu
 20

<210> 12
 <211> 24
 <212> PRT
 <213> Homo sapiens

<400> 12

Met Ser His Phe Asn Leu Tyr Leu Ile Leu Ile Met Cys Asn Ser Ile
 1 5 10 15

Ile Asp Pro Leu Ile Tyr Ala Leu
 20

<210> 13
 <211> 24
 <212> PRT
 <213> Rattus norvegicus

<400> 13

Met Ser His Phe Asn Leu Tyr Leu Ile Leu Ile Met Cys Asn Ala Val
 1 5 10 15

Ile Asp Pro Leu Ile Tyr Ala Leu
 20

<210> 14

<211> 23

<212> PRT

<213> Homo sapiens

<400> 14

Met Ser His Phe Asn Met Tyr Leu Ile Leu Ile Met Cys Asn Ser Val
 1 5 10 15

Ile Asp Pro Leu Ile Tyr Ala
 20

<210> 15

<211> 23

<212> PRT

<213> Mammalia sp.

<400> 15

Met Ser His Phe Asn Met Tyr Leu Ile Leu Ile Met Cys Asn Ser Val
 1 5 10 15

Ile Asp Pro Leu Ile Tyr Ala
 20

<210> 16

<211> 24

<212> PRT

<213> Homo sapiens

<400> 16

Met Ser Leu Phe Gln Val Asn Gly Val Leu Ile Met Cys Asn Ala Ile
 1 5 10 15

Ile Asp Pro Phe Ile Tyr Ala Leu
 20

<210> 17

<211> 22

<212> PRT

<213> Bos taurus

<400> 17

Ala His Phe Asn Thr Tyr Leu Val Leu Ile Met Cys Asn Ser Val Ile
 1 5 10 15

Asp Pro Leu Ile Tyr Ala

20

<210> 18
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 18

Ala His Phe Asn Thr Tyr Leu Val Leu Ile Met Cys Asn Ser Val Ile
 1 5 10 15

Asp Pro Leu Ile Tyr Ala
 20

<210> 19
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 19

Met Ser His Phe Asn Met Tyr Leu Ile Leu Ile Met Cys Asn Ser Val
 1 5 10 15

Met Asp Pro Leu Ile Tyr Ala
 20

<210> 20
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 20

Ser Tyr Phe Asn Leu Phe Leu Ile Leu Ile Ile Cys Asn Ser Val Val
 1 5 10 15

Asp Pro Leu Ile Tyr Ala
 20

<210> 21
 <211> 25
 <212> PRT
 <213> Bos taurus

<400> 21

Leu Ala Tyr Glu Lys Phe Phe Leu Leu Leu Ala Glu Phe Asn Ser Ala
 1 5 10 15

Met Asn Pro Ile Ile Tyr Ser Tyr Arg
 20 25

<210> 22
 <211> 19
 <212> PRT
 <213> Homo sapiens

<400> 22

Phe Leu Leu Leu Ala Glu Ala Asn Ser Leu Val Asn Ala Ala Val Tyr
1 5 10 15

Ser Cys Arg

<210> 23

<211> 22

<212> PRT

<213> Homo sapiens

<400> 23

Val Phe Ala Phe Cys Ser Met Leu Cys Leu Leu Asn Ser Thr Val Asn
1 5 10 15

Pro Leu Ile Tyr Ala Leu
20

<210> 24

<211> 21

<212> PRT

<213> Homo sapiens

<400> 24

Phe Gln Phe Phe Phe Trp Ile Gly Tyr Cys Asn Ser Ser Leu Asn Pro
1 5 10 15

Val Ile Tyr Thr Ile
20

<210> 25

<211> 22

<212> PRT

<213> Rattus norvegicus

<400> 25

Phe Asp Phe Val Val Ile Leu Thr Tyr Ala Asn Ser Cys Ala Asn Pro
1 5 10 15

Ile Leu Tyr Ala Phe Leu
20

<210> 26

<211> 16

<212> PRT

<213> Homo sapiens

<400> 26

Leu Ala Tyr Ser Asn Ser Ser Val Asn Pro Ile Ile Tyr Ala Phe Leu
1 5 10 15

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 October 2001 (11.10.2001)

PCT

(10) International Publication Number
WO 01/075161 A3

- (51) International Patent Classification⁷: **C12Q 1/68**
- (21) International Application Number: **PCT/US01/10076**
- (22) International Filing Date: **29 March 2001 (29.03.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
09/538,165 **30 March 2000 (30.03.2000)** **US**
- (71) Applicant (for all designated States except US): **IOWA STATE UNIVERSITY RESEARCH FOUNDATION** [US/US]; 310 Lab of Mechanics, Ames, IA 50011 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ROTHSCHILD, Max, F.** [US/US]; 310 Lab of Mechanics, Ames, IA 50011-2131 (US). **EMNETT, Rebecca** [US/US]; 310 Lab of Mechanics, Ames, IA 50011 (US). **KIM, Kwan, Suk** [US/US]; 310 Lab of Mechanics, Ames, IA 50011 (US).
- (74) Agent: **NEBEL, Heidi, S.**; Zarley, McKee, Thomte, Voorhees & Sease, Suite 3200, 801 Grand Avenue, Des Moines, IA 50309-2721 (US).
- (81) Designated States (national): **AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), **Eurasian** patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), **European** patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), **OAPI** patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report

[Continued on next page]

(54) Title: **GENETIC MARKERS FOR IMPROVED MEAT CHARACTERISTICS IN ANIMALS**

1 ACAAGAATCT GCATTCACCC ATGTACTTTT TCATCTGTAG CCTGGCTGTG
51 GCTGATATGC TGGTGAGCGT TTCCAATGGG TCAGAAACCA TTGTCATCAC
101 CCTATTAAAC AGCACGGACA CGGACGCACA GAGTTTCACA GTGAATATTG
151 ATAATGTCAT TGA CT CAGTG ATCTGTAGCT CCTTACTCGC CTCAATTTGC
201 AGCCTGCTTT CGATTGCAGT GGACAGGTAT TTTACTATCT TTTATGCTCT
251 CCAGTACCAT AACATTATGA CAGTTAAGCG GGTGGAATC ATCATCAGTT
301 GTATCTGGGC AGTCTGCACG GTGTCGGGTG TTTTGTTTCAT CATTTACTCA
351 GATAGCAGTG CTGTTATTAT CTGCCTCATA ACCGTGTTCT TCACCATGCT
401 GGCTCTCATG GCTTCTCTCT ATGTCCACAT GTTCCTCATG GCCAGACTCC
451 ACATTAAAGAG GATCGCCGTC CTCCCAGGCA CTGGCACCAT CCGCCAAGGT
501 GCCAACATGA AGGGGGCAAT TACCCTGACC ATCTTGATTG GGGTCTTTGT
551 GGTCTGCTGG GCCCCCTTCT TCCTCCACTT AATATTCTAT ATCTCCTGCC

(57) Abstract: Genetic markers in the porcine melanocortin-4 receptor (MC4R) gene are disclosed which are associated with favorable meat quality traits including, drip loss, marbling, pH and color. Further, novel sequence data from regions of the gene disclosed which may be used in a PCR test screen for the presence of the marker. The genetic marker may be used to screen animals for breeding purposes which have the desired traits. Kits which take advantage of the PCR test are also disclosed.

WO 01/075161 A3



(88) Date of publication of the international search report:
7 November 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/10076

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q G01N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, SEQUENCE SEARCH, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KIM K ET AL: "A missense variant of the porcine melanocortin-4 receptor (MC4R) gene is associated with fatness, growth and feed intake traits" MAMMALIAN GENOME, vol. 11, no. 2, February 2000 (2000-02), pages 131-135, XP002944421 the whole document ---	1-38
X	WO 00 06777 A (LARSON NIELS J ;UNIV IOWA RES FOUND (US); KIM KWAN SUK (US); ROTH S) 10 February 2000 (2000-02-10) the whole document --- -/--	1-39

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

8 July 2002

Date of mailing of the international search report

18/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Osborne, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/10076

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ANDERSSON L ET AL: "GENETIC MAPPING OF QUANTITATIVE TRAIT LOCI FOR GROWTH AND FATNESS IN PIGS" SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, US, vol. 263, 25 March 1994 (1994-03-25), pages 1771-1774, XP002018359 ISSN: 0036-8075 -----	
A	NEZER C ET AL: "An imprinted QLT with major effect on muscle mass and fat deposition maps to the IGF2 locus in pigs" NATURE GENETICS, vol. 21, no. 2, February 1999 (1999-02), pages 155-156, XP002204960 -----	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/10076

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 0006777	A	10-02-2000	AU	5230199 A		21-02-2000
			BR	9912460 A		13-02-2002
			CN	1328602 T		26-12-2001
			EP	1100970 A2		23-05-2001
			PL	346223 A1		28-01-2002
			WO	0006777 A2		10-02-2000
<hr/>						